


Many concepts in science involve applications of mathematics that measure certain quantities by their magnitude like length, mass, area, temperature, or energy. Only one number is needed to describe a length of 7 inches or 5°C for example. This single quantity is called **scalar**.

There are, however, many applications that involve not only the **magnitude** of an object but also, the **direction** of the displacement.

Vector: a quantity that has both magnitude and direction. For example, the flight pattern of a plane, has both speed (magnitude) and direction of travel. Velocity, acceleration, and force are described by both magnitude and direction and are known as vectors.




All vectors have two things:


- direction** – follow the arrow
- magnitude** – the length of the vector

I. Graphing Vectors


Use the vector to graph each of the following vectors.



1. $v - w$



2. $2v + 3w$



II. The Position Vector

To compute magnitude and direction of a vector, we need an algebraic way to describe the vector. The algebraic vector v is: $v = \langle a, b \rangle$

$$v = \langle \text{horizontal}, \text{vertical} \rangle$$

Where a and b are real (scalar) numbers and are called the components of the vector.

Vector v , may be described with initial point $P_1 (x_1, y_1)$ terminal point $P_2 (x_2, y_2)$

Vector v , is equal to the position vector: $v = \langle x_2 - x_1, y_2 - y_1 \rangle$

Find the position vector v with initial point $(-1, 2)$ and terminal point $(4, 6)$.

III. Vectors in terms of \mathbf{i} and \mathbf{j}

A vector of length **1** is called a **unit vector**. Let " \mathbf{i} " be a unit vector in the x-direction and " \mathbf{j} " be a unit vector in the y-direction. Any vector in the x-direction can be written as a scalar multiple of \mathbf{i} and any vector in the y-direction can be written as a scalar multiple of \mathbf{j} . They are defined as:

$$\mathbf{i} = \langle 1, 0 \rangle \text{ and } \mathbf{j} = \langle 0, 1 \rangle, \text{ where } \|\mathbf{i}\| = \sqrt{1^2 + 0^2} \text{ and } \|\mathbf{j}\| = \sqrt{0^2 + 1^2}.$$

$$\mathbf{v} = \langle a, b \rangle = a\langle 1, 0 \rangle + b\langle 0, 1 \rangle = a\mathbf{i} + b\mathbf{j}$$

Any vector may be expressed in terms of \mathbf{i} and \mathbf{j} .

A. Algebraic Operations

Vectors may be added, subtracted, or have scalar multiplication. Pretty straight forward, we can treat the numbers as coefficients and \mathbf{i} and \mathbf{j} as variables.

Let $\mathbf{v} = a_1\mathbf{i} + b_1\mathbf{j} = \langle a_1, b_1 \rangle$ and $\mathbf{w} = a_2\mathbf{i} + b_2\mathbf{j} = \langle a_2, b_2 \rangle$ be two vectors, and let α be a scalar. Then

$$\mathbf{v} + \mathbf{w} = (a_1 + a_2)\mathbf{i} + (b_1 + b_2)\mathbf{j} = \langle a_1 + a_2, b_1 + b_2 \rangle \quad (2)$$

$$\mathbf{v} - \mathbf{w} = (a_1 - a_2)\mathbf{i} + (b_1 - b_2)\mathbf{j} = \langle a_1 - a_2, b_1 - b_2 \rangle \quad (3)$$

$$\alpha\mathbf{v} = (\alpha a_1)\mathbf{i} + (\alpha b_1)\mathbf{j} = \langle \alpha a_1, \alpha b_1 \rangle \quad (4)$$

$$\|\mathbf{v}\| = \sqrt{a_1^2 + b_1^2} \quad (5)$$

If $\mathbf{v} = 2\mathbf{i} + 3\mathbf{j} = \langle 2, 3 \rangle$ and $\mathbf{w} = 3\mathbf{i} - 4\mathbf{j} = \langle 3, -4 \rangle$, find the following.

1. $\mathbf{v} + \mathbf{w}$

2. $\mathbf{v} - \mathbf{w}$

3. $3\mathbf{v}$

4. $2\mathbf{v} - 3\mathbf{w}$

5. $\|\mathbf{v}\|$

IV. Finding a vector from its Direction and Magnitude

Velocity vector - A vector that represents speed and direction of an object.

Force vector - A vector describing the direction and amount of force acting upon an object.

Given the magnitude $\|\mathbf{v}\|$ of a nonzero vector \mathbf{v} and the direction angle α , $0^\circ < \alpha < 360^\circ$, between vectors \mathbf{v} and \mathbf{i} , then: $\mathbf{v} = \|\mathbf{v}\|(\cos \alpha \mathbf{i} + \sin \alpha \mathbf{j})$

1. A ball is thrown with an initial speed of 25 mph in a direction that makes an angle of 30° with the positive x-axis. Express the velocity vector \mathbf{v} in terms of \mathbf{i} and \mathbf{j} . What is the initial speed in the horizontal direction? What is the initial speed in the vertical direction?

2. Find the direction angle α for $v = 4i - 4j$.

3. A Boeing 737 aircraft maintains a constant airspeed of 500 mph headed due south. The jet stream is 80 mph in the northeasterly direction.

a) Express the velocity v_a of the 737 relative to the air and velocity v_w of the jet stream in terms of i and j .

b) Find the velocity of the 737 relative to the ground.

c) Find the actual speed and direction of the 737 relative to the ground.

4. Two movers require a magnitude of force of 300 pounds to push a piano up a ramp inclined at an angle 20° from the horizontal. How much does the piano weigh?

An object in **Static Equilibrium**: the object is at rest and the sum of all forces acting on the object is zero, a.k.a. the resultant force is zero.

5. A box of supplies that weighs 1200 pounds is suspended by two cables attached to the ceiling. What are the tensions in the two cables?

